

GRAD-SHAFRANOV EQUATION WITH ANISOTROPIC PRESSURE

VASILY S. BESKIN

P.N. Lebedev Physical Institute, Russian Academy of Sciences, Moscow, 117924, Russia; beskin@lpi.ru
 DAEC, Observatory Paris – Meudon, 92125, Meudon, France

AND

INGA V. KUZNETSOVA

Moscow Institute of Physics and Technology, Dolgoprudnyi, 141700, Russia; kuzn@lpi.ru

submitted to ApJ, February 25, 2000

ABSTRACT

The most general form of the nonrelativistic Grad-Shafranov equation describing anisotropic pressure effects is formulated within the double adiabatic approximation. It gives a possibility to analyze quantitatively how the anisotropic pressure affects the 2D structure of the ideal magnetohydrodynamical flows.

Subject headings: MHD — star: winds, outflows — ISM: jets and outflows

1. INTRODUCTION

The Grad-Shafranov (GS) equation for axisymmetric stationary ideal magnetohydrodynamical (MHD) flows has been considered very intensively within the last two decades both for nonrelativistic stellar (solar) wind including outflow from young stellar objects (Okamoto 1975; Heinemann & Olbert 1978; Blandford & Payne 1982; Heyvaerts & Norman 1989; Sakurai 1990; Pelletier & Pudritz 1992; Shu et al 1994), and for relativistic outflows from radio pulsars (Ardavan 1979; Bogovalov 1992) and 'central engine' in active galactic nuclei (Phinney 1983; Begelman, Blandford & Rees 1984; Mobarry & Lovelace 1988; Chiu, Li & Begelman 1991; Beskin, Kuznetsova & Rafikov 1998). As is well known, in this approach the isotropic gas pressure has been postulated (see e.g. Heyvaerts 1996 for introduction).

It is necessary to stress that, actually, thermal effects were considered rather carefully only within the given poloidal magnetic field approximation (Weber & Davis 1967; Kennel, Fujimura & Okamoto 1983; Paatz & Camenzind 1996), or within the self-similar approach (Low & Tsinganos 1986; Contopoulos & Lovelace 1994; Sauty & Tsinganos 1994). As to the GS equation, its 2D solutions with nonzero pressure were only obtained for pure hydrodynamical flows (see Beskin 1997 for review) when the isotropic pressure model is correct.

On the other hand, for real astrophysical flows where the effects of finite pressure may play an important role, the isotropic pressure model does not seem to be sufficient. For example, in the solar wind where the free path l_ν exceeds the Larmor radius r_H by many order of magnitude, so that $l_\nu/r_H \sim 10^9$ in the vicinity of the Earth, the effect of the anisotropic pressure may be important.

Of course, it is necessary to remember that the ideal MHD approximation (including the double adiabatic approach under consideration) is not good for real astrophysical objects, and so the dissipative processes can play an important role (Leer & Axford 1972; Hu, Esser & Habbal 1997). Moreover, in the solar wind the stochastic component of the magnetic field is of the order of the regular one. Such fluctuations drastically change the velocity dis-

tribution function. As a result, the pressure anisotropy $P_\parallel/P_\perp \approx 1.2$ is not actually large (see e.g. Marsch et al 1982). May be, for this reason the effects of the anisotropic pressure on the outflow structure within the GS approach were analyzed in a small number of papers (Asséo & Beaulieu 1983; Tsikarishvili, Rogava & Tsikauri 1995), the GS equation itself being formulated only for the planar static configurations with zero longitudinal electric current (Nötzel, Schindler & Birn 1985).

In this paper we give the general equations describing the axisymmetric stationary ideal MHD flows with anisotropic pressure. Then, we shall present the results of our preliminary analysis of the effects of the anisotropic pressure (such an analysis has not been produced previously either).

2. BASIC EQUATIONS

Let us consider a stationary axisymmetric (and nonrelativistic) MHD flow. Then the magnetic field can be written in the form

$$\mathbf{B} = \frac{\nabla \Psi \times \mathbf{e}_{\hat{\varphi}}}{2\pi\varpi} - \frac{2I}{\varpi c} \mathbf{e}_{\hat{\varphi}}. \quad (1)$$

Here $\Psi(r, \theta)$ is the magnetic flux, $\varpi = r \sin \theta$, and $I(r, \theta)$ is the total electric current flowing inside the region $\Psi < \Psi(r, \theta)$. Assuming that the magnetosphere contains enough plasma to screen the longitudinal electric field $E_\parallel$, one can write down

$$\mathbf{E} = -\frac{\Omega_F}{2\pi c} \nabla \Psi, \quad (2)$$

the 'field angular velocity' $\Omega_F = \Omega_F(\Psi)$ being constant on the magnetic surfaces $\Psi(r, \theta) = \text{const}$. Finally, the continuity equation $\nabla \cdot (n\mathbf{u}) = 0$ together with the freezing-in condition $\mathbf{E} + \mathbf{v} \times \mathbf{B}/c = 0$ and the Maxwell equation $\nabla \times \mathbf{E} = 0$ result in

$$\mathbf{v} = \frac{\eta}{\rho} \mathbf{B} + \Omega_F \varpi \mathbf{e}_{\hat{\varphi}}, \quad (3)$$

where ρ is the mass density and $\eta = \eta(\Psi)$ is the particle-to-magnetic flux ratio. Clearly, these expressions do not depend on the pressure anisotropy.

The next step is the transformation of the Euler equation

$$\rho(\mathbf{v} \cdot \nabla)\mathbf{v} = -\nabla_\beta P_{\alpha\beta} + \frac{1}{4\pi}[\nabla \times \mathbf{B}] \times \mathbf{B} + \rho\mathbf{g}. \quad (4)$$

According to Chew, Goldberger & Low (1956), for collisionless plasma with an anisotropic pressure

$$P_{\alpha\beta} = P_s \delta_{\alpha\beta} + (P_n - P_s) B_\alpha B_\beta / B^2 \quad (5)$$

it is possible to introduce two extra invariants on the magnetic surfaces $\Psi(r, \theta) = \text{const}$ (the so-called double adiabatic approximation resulting from the conservation of two adiabatic invariants)

$$s_1(\Psi) = \frac{P_n B^2}{\rho^3}, \quad (6)$$

$$s_2(\Psi) = \frac{P_s}{\rho B}. \quad (7)$$

They correspond to polytropic equations of state with $\Gamma_\parallel = 3$ and $\Gamma_\perp = 1$. As a result, integrating $\mathbf{v}$ and φ components of equation (4), one can obtain the known expressions for the energy and the angular momentum (Asséo & Beaufils 1983; Tsikarishvili, Rogava & Tsikauri 1995):

$$E(\Psi) = \frac{v^2}{2} + \frac{P_s}{\rho} + \frac{3}{2} \frac{P_n}{\rho} + \frac{\Omega_F I}{2\pi c\eta} (1 - \beta_a) - \frac{GM}{r}, \quad (8)$$

$$L(\Psi) = v_\varphi r \sin \theta + \frac{I}{2\pi c\eta} (1 - \beta_a). \quad (9)$$

Here

$$\beta_a = 4\pi \frac{P_n - P_s}{B^2} \quad (10)$$

is the anisotropic pressure parameter. Together with $\Omega_F(\Psi)$ and $\eta(\Psi)$ these six invariants determine all the characteristics of a flow. In particular,

$$\frac{I}{2\pi} = c\eta \frac{L - \Omega_F \varpi^2}{1 - M^2 - \beta_a}, \quad (11)$$

$$v_\varphi = \frac{1}{\varpi} \cdot \frac{\Omega_F \varpi^2 (1 - \beta_a) - M^2 L}{1 - M^2 - \beta_a}, \quad (12)$$

where $M^2 = 4\pi\eta^2/\rho$ is the Alfvénic Mach number. Thus, in the anisotropic case the Alfvénic singularity has the form

$$A = 1 - M^2 - \beta_a = 0. \quad (13)$$

It is necessary to stress that (11) gives an implicit expression for the current I because I through B_φ^2 is contained in the rhs of this equation as well. Nevertheless, the standard procedure of the determination of flow parameters in a given poloidal magnetic field $\mathbf{B}_p$ remains the same. The Bernoulli equation (8), which can be rewritten in the form

$$\begin{aligned} \frac{M^4}{64\pi^4\eta^2} (\nabla\Psi)^2 &= 2\varpi^2 \left(E - \frac{P_s}{\rho} - \frac{3}{2} \frac{P_n}{\rho} + \frac{GM}{r} \right) \\ &\quad - \frac{[\Omega_F \varpi^2 (1 - \beta_a) - LM^2]^2}{(1 - M^2 - \beta_a)^2} \\ &\quad - 2\varpi^2 \Omega_F (1 - \beta_a) \frac{L - \Omega_F \varpi^2}{1 - M^2 - \beta_a}, \end{aligned} \quad (14)$$

taken together with (11) and with the definitions (6), (7) determines implicitly the Mach number M^2 and β_a as functions of Ψ (or $\mathbf{B}_p$) and all the six invariants

$$M^2 = M^2[(\nabla\Psi)^2, E, L, \Omega_F, \eta, s_1, s_2], \quad (15)$$

$$\beta_a = \beta_a[(\nabla\Psi)^2, E, L, \Omega_F, \eta, s_1, s_2]. \quad (16)$$

It means that all the other parameters (I , v_φ , etc.) can be found from given values of $\mathbf{B}_p$ and of all the six invariants as well. This fact is fully analogous to spherically symmetric flows (Bondi 1952; Parker 1958) where the flow structure is known and the algebraic expressions for the integrals of motion are enough to determine all the characteristics of a flow.

On the other hand, if the flow is not spherically symmetric, the poloidal magnetic field is to be determined from the $\nabla\Psi$ component of equation (4). After numerous but elementary calculations it can be presented in the following compact form

$$\begin{aligned} \nabla_k \left[\frac{1}{\varpi^2} (1 - M^2 - \beta_a) \nabla^k \Psi \right] &+ \frac{64\pi^4}{\varpi^2} \frac{1}{2M^2} \frac{\partial}{\partial \Psi} \left(\frac{G}{A} \right) \\ &- 8\pi^3 P_n \frac{1}{s_1} \frac{ds_1}{d\Psi} - 16\pi^3 P_s \frac{1}{s_2} \frac{ds_2}{d\Psi} = 0 \end{aligned} \quad (17)$$

Here

$$\begin{aligned} \left(\frac{G}{A} \right) &= 2\varpi^2 \eta^2 \left(E - \frac{P_s}{\rho} - \frac{3}{2} \frac{P_n}{\rho} + \frac{GM}{r} \right) \\ &+ \eta^2 \frac{\varpi^4 \Omega_F^2 (1 - \beta_a) - 2\varpi^2 \Omega_F L (1 - \beta_a) + M^2 L^2}{1 - M^2 - \beta_a}, \end{aligned} \quad (18)$$

∇_k is a covariant derivative, and the operator $\partial/\partial\Psi$ acts on the invariants $E(\Psi)$, $L(\Psi)$, $\Omega_F(\Psi)$, and $\eta(\Psi)$ only. According to (6)–(7) and (15)–(16), this trans-field (stream, generalized GS) equation contains the unknown function $\Psi(r, \theta)$ and the six invariants only. As to the GS equation itself describing statical configurations, it can be obtained from (17) in the limit $\Omega_F \rightarrow 0$, $\eta \rightarrow 0$ (and hence $M^2 \rightarrow 0$), but $\eta L \rightarrow \text{const}$. It gives for gravity-free case

$$\begin{aligned} \nabla_k \left[\frac{1}{\varpi^2} (1 - \beta_a) \nabla^k \Psi \right] &+ \frac{16\pi^2}{\varpi^2} (1 - \beta_a) I \frac{dI}{d\Psi} \\ &+ 16\pi^3 \rho \frac{d}{d\Psi} \left(\frac{P_s}{\rho} + \frac{3}{2} \frac{P_n}{\rho} \right) \\ &- 8\pi^3 P_n \frac{1}{s_1} \frac{ds_1}{d\Psi} - 16\pi^3 P_s \frac{1}{s_2} \frac{ds_2}{d\Psi} = 0. \end{aligned} \quad (19)$$

Relations (8)–(12) remain true even for a more general bounded anisotropy model (Denton et al 1994)

$$(\mathbf{v} \cdot \nabla) \left(\frac{P_n B^2}{\rho^3} \right) = 2\mathcal{P}_t \left(\frac{B^2}{\rho^3} \right), \quad (20)$$

$$(\mathbf{v} \cdot \nabla) \left(\frac{P_s}{\rho B} \right) = -\mathcal{P}_t \left(\frac{1}{\rho B} \right), \quad (21)$$

where $\mathcal{P}_t$ is a nondissipative energy exchange term. In this case we have the conservation of the ‘total entropy’ $S = S(\Psi)$

$$(\mathbf{v} \cdot \nabla) S = \frac{1}{2} \frac{\rho^3}{B^2} (\mathbf{v} \cdot \nabla) s_1 + \rho B (\mathbf{v} \cdot \nabla) s_2 = 0, \quad (22)$$

and so the generalized GS equation (17) has a form

$$\nabla_k \left[\frac{1}{\varpi^2} (1 - M^2 - \beta_a) \nabla^k \Psi \right] + \frac{32\pi^4}{\varpi^2 M^2} \frac{\partial}{\partial \Psi} \left(\frac{G}{A} \right) - 16\pi^3 \frac{dS}{d\Psi} = 0. \quad (23)$$

Introducing the effective pressure $P_{\text{eff}} = P_{\text{eff}}(\Psi)$ as

$$dP_{\text{eff}} = \rho d \left(\frac{P_s}{\rho} + \frac{3}{2} \frac{P_n}{\rho} \right) - dS, \quad (24)$$

which is equivalent to the ordinary thermodynamic relation $dP = \rho dw - \rho T ds$, one can obtain instead of (19)

$$\nabla_k \left[\frac{1}{\varpi^2} (1 - \beta_a) \nabla^k \Psi \right] + \frac{16\pi^2}{\varpi^2} (1 - \beta_a) I \frac{dI}{d\Psi} + 16\pi^3 \frac{dP_{\text{eff}}}{d\Psi} = 0. \quad (25)$$

For $I = 0$ it has the form found by Nötzel, Schindler & Birn (1985) for planar geometry.

Equation (17) is a partial equation of the mixed type. Using the implicit relations resulting in (15)–(16), one can find for the second-order operator in (17)

$$A \left[\nabla_k \left(\frac{1}{\varpi^2} \nabla^k \Psi \right) + \frac{(\nabla^i \Psi)(\nabla^k \Psi) \nabla_i \nabla_k \Psi}{\varpi^2 (\nabla \Psi)^2 D} \right] + \dots = 0, \quad (26)$$

where

$$D = \frac{n}{d}, \quad (27)$$

and

$$n = \left(1 - M^2 - \beta_a + 4\pi \frac{4P_n - P_s}{B^2} \frac{B_\varphi^2}{B^2} \right) \left(1 - 3 \frac{P_n}{\rho v_p^2} \right) + \frac{B_\varphi^2}{B_p^2} \left(1 - \beta_a - 4\pi \frac{3P_n - P_s}{B^2} + 4\pi \frac{4P_n - P_s}{B^2} \frac{B_\varphi^2}{B^2} \right) - \left(M^2 - 4\pi \frac{3P_n - P_s}{B^2} \right) \frac{3P_n - P_s}{\rho v_p^2} \frac{B_\varphi^2}{B^2} \quad (28)$$

$$d = \left(1 - 3 \frac{P_n}{\rho v_p^2} \frac{B_p^2}{B^2} \right) \left(M^2 + 4\pi \frac{P_n}{B^2} \right) + \frac{P_s}{\rho v_p^2} \frac{B_p^2}{B^2} \left(M^2 - 4\pi \frac{3P_n - P_s}{B^2} \right). \quad (29)$$

The form (26) of equation (17) coincides with the one found for an isotropic pressure (see e.g. Sakurai 1990; Beskin 1997). Hence, one can conclude that the trans-field equation is elliptical for $D > 0$ and $D < -1$, and hyperbolic for $-1 < D < 0$. It means that the trans-field equation changes from elliptical to hyperbolic on the surfaces $D = 0$ (fast and slow magnetosonic singularities) and $D = -1$ (cusp singularity).

It is necessary to stress that, as in the isotropic case, D can be presented as $D = d_1 A + d_2 B_\varphi^2/B^2$, so that $D \propto A$ for $B_\varphi = 0$. This property reflects the fact that for $\mathbf{k} \parallel \mathbf{B}$ fast or slow magnetosonic velocity coincides with

the Alfvénic one. In particular, for a nonrotating flow ($\Omega_F = 0$, $L = 0$), when, according to (11), $B_\varphi = 0$, we have

$$n = (1 - M^2 - \beta_a) \left(1 - 3 \frac{P_n}{\rho v_p^2} \right). \quad (30)$$

Using now the definitions (13) and (27), one can obtain for sonic, Alfvénic, and cusp velocities in anisotropic media

$$a_s^2 = 3 \frac{P_n}{\rho}, \quad (31)$$

$$v_A^2 = \frac{B^2}{4\pi\rho} - \frac{P_n - P_s}{\rho}, \quad (32)$$

$$v_c^2 = \frac{3 \frac{P_n}{\rho} \frac{B^2}{4\pi\rho} + 6 \frac{P_n P_s}{\rho^2} - \frac{P_s^2}{\rho^2}}{\frac{B^2}{4\pi\rho} + 2 \frac{P_s}{\rho}}. \quad (33)$$

As we see, the velocities (31)–(33) coincide exactly with the characteristic velocities propagating in anisotropic media (Clemmow & Dougherty 1969). Recall that the condition $v_A^2 < 0$ corresponds to the firehose instability, and the condition $v_c^2 < 0$ – to the mirror instability.

3. TWO EXAMPLES

3.1. Free outflow

As a first example, let us consider a free (i.e. without gravity effects) outflow along a strong monopole magnetic field from the surface of a nonrotating sphere (with radius r_0 , magnetic field B_0 , pressure $P_0 \ll B_0^2/4\pi$, mass density ρ_0 , and radial velocity v_0). It is natural to assume that here $P_s(r_0) = P_n(r_0) = P_0$. As a result, one can find for the radial velocity

$$\frac{v^2}{v_0^2} = \frac{1}{2} + \frac{5}{2} p \left(1 - \frac{2}{5} \frac{r_0^2}{r^2} \right) + \frac{1}{2} \sqrt{\left[1 + 5p \left(1 - \frac{2}{5} \frac{r_0^2}{r^2} \right) \right]^2 - 12p}, \quad (34)$$

where

$$p = \frac{P_0}{\rho_0 v_0^2}. \quad (35)$$

It gives

$$r \frac{d(v^2/v_0^2)}{dr} = \frac{N_v}{D}, \quad (36)$$

where $N_v = 4pr_0^2/r^2$, and

$$D = 1 - 3 \frac{P_n}{\rho v^2}, \quad (37)$$

which corresponds to the longitudinal sonic singularity in (27). On the other hand, the same as for isotropic media, for a nonrotating wind the magnetosonic surface coincides with the Alfvénic one, the singularity for $1 - M^2 - \beta_a = 0$ being absent.

Comparing now expression (34) for $v(r)$ with the appropriate one corresponding to an outflow with an isotropic pressure for the polytropic equation of state with $\Gamma = 3$

and the same energy E , one can conclude that the difference only takes place for small enough pressure $P_0 \ll \rho_0 v_0^2$ when

$$\frac{v_{\infty}^{(\text{iso})}}{v_0} = 1 + \frac{1}{6} \left(\frac{2E}{v_0^2} - 1 \right), \quad (38)$$

$$\frac{v_{\infty}^{(\text{aniso})}}{v_0} = 1 + \frac{1}{5} \left(\frac{2E}{v_0^2} - 1 \right). \quad (39)$$

For $P_0 \gg \rho_0 v_0^2$ the limiting velocity does not depend on the pressure model: $v_{\infty}^2 \approx 2E$. Thus, the difference takes place for $r \sim r_0$ only.

It is also interesting to determine how the anisotropic pressure affects the position of the Alfvénic surface r_{cr} coinciding for $\Omega_F = 0$ with the fast magnetosonic one. As

$$\frac{\beta_a}{M^2} \approx p \left(\frac{v_0}{v} \right)^4, \quad (40)$$

it is easily check that $\beta_a/M^2 \ll 1$ for both $p \ll 1$ and $p \gg 1$, so that for $P_0 \ll \rho_0 v_0^2$ and $P_0 \gg \rho_0 v_0^2$ the position of the Alfvénic surface

$$r_{\text{cr}}^2 \approx r_0^2 \frac{B_0^2}{4\pi\rho_0 v_0^2} \frac{v_0}{v_{\infty}} \quad (41)$$

does not depend on the pressure anisotropy. Finally, as

$$\frac{P_n}{P_s} = \frac{v_0^2}{v^2} \frac{r^2}{r_0^2}, \quad (42)$$

the anisotropy of the pressure tensor increases with distance from the central object. As has already been stressed, it can be realized only if dissipative processes (heat conductivity, plasma instability, etc.) play no role. Relations (40) and (42) also mean that the conditions for the firehose and mirror instabilities are not satisfied up to the Alfvénic surface. On the other hand, the firehose instability takes place in the supersonic region for $r > r_{\text{fh}} \gg r_{\text{cr}}$, where

$$r_{\text{fh}}^2 \approx r_{\text{cr}}^2 \frac{1}{p} \left(\frac{v_{\infty}}{v_0} \right)^4. \quad (43)$$

3.2. Parker outflow

As a next example we consider an outflow from the gravity center of mass M in a monopole magnetic field (Parker ejection). Analyzing the energy equation (8), we have

$$\frac{v^2}{v_0^2} = \frac{1}{2} + \frac{5}{2}p - p \frac{r_0^2}{r^2} + \frac{GM}{v_0^2} \frac{r_0 - r}{rr_0} + \frac{1}{2} \sqrt{\left(1 + 5p - 2p \frac{r_0^2}{r^2} + 2 \frac{GM}{v_0^2} \frac{r_0 - r}{rr_0} \right)^2 - 12p}. \quad (44)$$

As a result, now $N_v = 4pr_0^2/r^2 - 2GM/v_0^2 r$, and D is determined by (37) again. Hence, for the position of the sonic surface we have

$$r_s = 2r_0 \frac{P_0 r_0}{GM \rho_0}. \quad (45)$$

On the other hand, one can easily check that equation (44) has a real solution for $r_s \leq r_0$ only. This result is in agreement with a well-known fact that no transonic outflow is possible for a gas with a polytropic index $\Gamma = 3$. Simultaneously, for Parker ejection the conditions of the firehose and mirror instabilities up to Alfvénic surface are not satisfied either.

4. CONCLUSION

Thus, we have generalized the GS equation for the flows with anisotropic pressure. It allows a quantitative analysis of the effect of anisotropic pressure upon the 2D outflow structure within ideal double adiabatic approximation. As has been shown on simple examples, the pressure anisotropy itself may be very large, but not in all cases does it result in a large disturbance of the flow parameters. Clearly, the additional analysis (say, within the Weber-Davis model) is necessary to clarify all the details of the flow with anisotropic pressure.

This work was supported by the Russian Foundation for Basic Research (project No. 99-02-17184). VSB thanks A.A. Samsonov for stimulating discussions and Observatory Paris – Meudon for hospitality. IVK thanks the International Soros Educational Program (Grant a99-941) and Landau Foundation for financial support.

REFERENCES

- Ardavan, H. 1979, MNRAS, 189, 397
 Asséo, E., & Beaufils, D. 1983, Ap&SS, 89, 133
 Begelman, M.C., Blandford, R.D., & Rees, M.J. 1984, Rev. Mod. Phys., 56, 255
 Beskin, V.S. 1997, Physics Uspekhi, 40(7), 659
 Beskin, V.S., Kuznetsova, I.V., & Rafikov, R.R. 1998, MNRAS, 299, 341
 Blandford, R.D., & Payne, D.G. 1982, MNRAS, 199, 883
 Bogovalov, S.V. 1992, Sov. Astron. Lett., 18, 337
 Bondi, H. 1952, MNRAS, 112, 195
 Chew, G.F., Goldberger, M.L., & Low, F.E. 1956, Proc. Roy. Soc. London A, 236, 112
 Chiueh, T., Li, Zh.-Y., & Begelman, M.C. 1991, ApJ, 377, 462
 Clemmow, P.C., & Dougherty, J.P. 1969, Electrodynamics of Particles and Plasmas, (Reading: Addison-Wesley Publ. C.), 385
 Contopoulos, J., & Lovelace, R.V.E. 1994, ApJ, 429, 139
 Denton, R.E., Anderson, B.J., Gary, S.P., & Fuselier, S.A. 1994, J. Geophys. Res., 99, 11225
 Heinemann, M., & Olbert, S. 1978, J. Geophys. Res., 83, 2457
 Heyvaerts, J., 1996, in Plasma Astrophysics, ed. C. Chiuderi & G. Einaudi (Berlin: Springer), 31
 Heyvaerts, J., & Norman, C. 1989, ApJ, 347, 1055
 Hu, Y.Q., Esser, R., & Habbal, S.R. 1997, J. Geophys. Res., 102, 14661
 Kennel, C.F., Fujimura, F.S., & Okamoto, I. 1983, Geophys. Astrophys. Fluid Dynamics, 26, 147
 Leer, L., & Axford, W.I. 1972, Sol. Phys., 23, 238
 Low, B.C., & Tsinganos, K. 1986, ApJ, 302, 163
 Marsch, E., et al. 1982, J. Geophys. Res., 87, 52
 Mobarri, C.M., & Lovelace, R.V.E. 1986, ApJ, 309, 455
 Nötzel, A., Schindler, K., & Birn, J. 1985, J. Geophys. Res., 90, 8293
 Okamoto, I. 1975, MNRAS, 173, 357
 Paatz, G., & Camenzind, M. 1996, A&A, 313, 591
 Parker, E.N. 1958, ApJ, 128, 664
 Pelletier, G., & Pudritz, R.E. 1992, ApJ, 394, 117
 Phinney, E. 1983, in Astrophysical Jets, ed. A. Ferrary & A.G. Pacholczyk, (Dordrecht: Reidel), 201
 Sakurai, T. 1990, Computer Phys. Rep., 12, 247
 Sauty, C., & Tsinganos, K. 1994, A&A, 287, 893
 Shu, F.H., et al. 1994, ApJ, 429, 797
 Tsikarishvili, E.G., Rogava, A.D., & Tsikauri, D.G. 1995, ApJ, 439, 822
 Weber, E.J., & Davis, L. Jr 1967, ApJ, 148, 217